

# FRIDAY MORNING SESSION

December 1, 1967

The Friday morning session convened at 9:00 a. m. in Ballroom A of the Admiral Semmes Hotel. Dr. Thomas F. Cannon served as moderator.

MODERATOR CANNON: I think we have a very excellent program this morning and to lead it off we have Dr. Ronald M. Girouard who will speak on the "Anatomy of Adventitious Root Formation".

## ANATOMY OF ADVENTITIOUS ROOT FORMATION IN STEM CUTTINGS

RONALD M. GIROUARD

*Forest Research Laboratory,  
Department of Forestry and Rural Development,  
Sillery, Quebec, Canada*

The art of propagating plants from cuttings is old and may have started with man's initial interest in ornamental, medicinal, orchard and field plants. But as a science, a series of facts systematically arranged to explain or predict the operation of general laws, the propagation of plants from cuttings did not begin to develop until the eighteenth century (24). Probably the first paper describing the scientific approach to plant propagation was written by the French dendrologist Duhamel du Monceau in 1758 (see fig. 1 and ref. 10). My paper this morning will be an attempt to summarize what is known concerning the anatomy of adventitious root formation in stem cuttings. And when compared with the volumes written by Duhamel, my talk will simply resemble a drop in the bucket.

The rooting of stem cuttings is one of the widely used forms of asexual propagation, a fact which you all know. It involves the placing of stem pieces with one or more buds under conditions that favor the regeneration of roots. Stem cuttings we generally classify as hardwood, softwood and herbaceous depending upon the degree of lignification of the stems, the presence or absence of leaves, and the plant species involved.

Plant propagators are aware of the fact that several easy-to-root plants bear aerial roots or roots in early stages of development within the tissues of the bark. This last group has been referred to by some authors as preformed, hidden, latent and initial roots or root germs. The term morphological roots, first used by the Dutch investigator van der Lek (37), designates all of the roots present when the stems are cut from the stock plants. Figures 2 and 3 show aerial roots that have form-




Figure 1. As early as 1758 efforts were made to study plant propagation from a scientific viewpoint. Girdling, grafting, polarity of shoots used as cuttings, and layering were a few of the subjects studied. Redrawn from H. L. Duhamel du Monceau. *La physique des arbres*. Guerin & Delatour, Paris. Plate XV, volume 2. 1758.


ed just below a node on stems taken from a juvenile plant of *Hedera helix*. In figure 3 the stem was stained with basic fuchsin, cleared and photographed while in methyl salicylate; the vascular bundles have retained the stain. A list of woody plants with stems bearing aerial and latent roots (morphological roots) is given in table 1.

The most common adventitious roots of stems are those that form completely after the cuttings are made; they develop near the base of stems from wounded tissues or above the cuts. It is not surprising that these structures have been named wound or wound-induced roots (18,22,37).

Since most adventitious roots arise endogenously, that is, in tissues located at some distance from the periphery of the stems, it is only appropriate that we should be familiar with some features of the internal structure of stems. First, let us consider the subdivisions of the vascular system, the vascular bundles or fascicles (traces) which function as the transport network of young shoots.

In stems of conifers and woody dicotyledons, the internodes have vascular bundles arranged in rings but separated by panels of cells high in metabolic activity. In cross-sections of stems of a number of herbaceous dicotyledons and many

monocotyledons, the vascular fascicles appear scattered. Figure 4 gives us a general idea of the distribution of the bundles as found in several of the plants that we propagate. A closer look at one of the vascular bundles of a woody dicotyledon (fig. 5) reveals two smaller strands or traces: one composed of xylem



Figures 2 and 3. Aerial roots that have formed below a node on stems taken from a juvenile plant of *Hedera helix* are readily seen. In figure 3, the stem was stained with basic fuchsin, cleared and photographed while in methyl salicylate, a clearing agent. The development of vascular traces into the new roots is noted. Figure 3: from Girouard, R. M. *Canad. J. Bot.* 45:1877-1881.

Table 1. Woody plants with stems bearing aerial and latent roots. \*

<i>Campsis radicans</i>	<i>Populus nigra thevestina</i> (37)
<i>Cotoneaster dammeri</i> (38)	<i>Populus simonii</i> (37)
<i>Eugeissona</i> spp. (35)	<i>Populus trichocarpa</i> (2,3,37)
<i>Ficus carica</i> (11)	<i>Ribes alpinum</i> (37)
<i>Hedera helix</i> (36)	<i>Salix alba</i> (36)
<i>Jasminum</i> spp. (18)	<i>Salix alba chermenisa</i> (37)
<i>Lonicera japonica</i> (27)	<i>Salix alba sericea</i> (37)
<i>Maricaria</i> spp. (35)	<i>Salix amygdalina</i> (37)
<i>Populus</i> Section <i>Aegeiros</i> (4)	<i>Salix babylonica</i> (37)
<i>Populus alba</i> (25)	<i>Salix discolor</i> (22,37)
<i>Populus x canadensis</i> (2,3)	<i>Salix fragilis</i> (6,7)
<i>Populus deltoides</i> (25)	<i>Salix myrsinifolia</i> (37)
<i>Populus x euramericana</i> (25)	<i>Salix viminalis</i> (36)
<i>Populus nigra italica</i>	<i>Vershaeffeltia</i> spp. (35)
(29,36,37)	

\* Numbers refer to references cited.




Figure 4. The vascular bundles of conifers and woody dicotyledons at the internodes are arranged in rings (A, B). In cross-sections of stems of a number of herbaceous dicotyledons and many monocotyledons the vascular fascicles are scattered (C,D). Fig. 4 A, C From: B. Huber, 1961, *Grundzuge der Pflanzenanatomie*. Springer-Verlag, Berlin. p. 84.

(wood) and another of phloem elements. Between the strands of xylem and phloem are cells of a lateral meristem — the fascicular cambium. At a later stage, cambial cells differentiate between the bundles to form the interfascicular cambium. It is the vascular cambium made up of the fascicular and interfascicular cambia which is responsible for the secondary growth of stems. Much of this growth is readily identified as annual growth rings in the secondary xylem. Now for a few minutes let us consider the vascular system at the nodes.

The presence of a leaf results in one or more bundles bending away from the main formation of the vascular system and in connecting the leaf to the stem (fig. 6). Bundles of this kind can be traced down through the stem to a point where they join other strands. These links between the leaf and the stem are the leaf traces. Immediately above the bending part of the

traces in the stem, parenchyma cells with unspecialized properties contrast with the highly specialized cells of the vascular tissues. These groups of parenchyma cells make up the leaf gaps. Buds that form in the axils of leaves are compressed shoots or miniature branches. Nevertheless they too are linked to the main stem but by branch traces; they also have branch gaps.

I am certain that many of you like myself have wondered just where in stems the adventitious roots originate and develop, and what anatomical factors influence their development. The answer to these questions have not always been easy to obtain and they have varied considerably from one plant species to the other. I will attempt to summarize studies that have come to my attention.

It is definitely known that adventitious roots have a tendency to arise in the vicinity of differentiating tissues of the vascular bundles; the proximity of the new roots to the main axis of the stems permits a rapid vascular connection between the plant parts. In young stems of dicotyledons and gymnosperms, roots are initiated by cells located near the periphery of the vascular fascicles, often interfascicular parenchyma cells, while in relatively old stems the seats of initiation are




Figure 5. A close look at a cross-section of a young woody dicotyledon shows vascular bundles arranged in a ring and individual bundles composed of two strands: one of xylem (wood) and another of phloem elements. E, epidermis; C, cortex; P, phloem; X, xylem; PI, pith; VB, vascular bundle; FC, fascicular cambium; IC, interfascicular cambium. From: Girouard, R. M. 1967. *Canad. J. Bot.* 45:1877-1881.




Figure 6 Leaves and buds are connected to the vascular system of stems by leaf traces and branch traces, respectively. Both leaves and buds have gaps made up of parenchyma cells above their respective traces. LT, leaf trace; LG, leaf gap; BT, branch trace. From K. Esau 1960 *Anatomy of seed plants*. John Wiley & Sons, New York, p. 209.

close to the interfascicular cambium. In monocotyledons, adventitious roots arise in parenchyma cells external to the vascular tissues. In every case cells that are alive, not just walls, and that are capable of dividing are necessary for the formation of adventitious organs such as roots.

These structures can arise internally (endogenously) not only in nodal tissues adjacent to leaf and branch gaps but in internodes. They can occur externally (exogenously) in callus tissue formed at the basal end of cuttings and in hyperhydric outgrowths of lenticels. Tables 2 and 3 provide information concerning the seat of origin of morphological and wound-induced roots in several woody plants.

Up to this point I have brought to your attention the vascular system at nodes and internodes and the sites of root initiation, but one important fact remains to be mentioned. It is this: while a root may be formed within tissues of a stem, its emergence is not a sure thing. A correlation between the presence of many thick-walled cells such as fibers and sclereids (two components of sclerenchyma) and poor rooting of shoots has been reported by several investigators (1,8,9,17,23). Some workers believe that the groups of fibers act as mechanical barriers preventing the lateral emergence of roots and promoting basal emergence through callus tissue. Recently a few scientists (15,16,23) have been unable to find simple relation-

Table 2 Origin of aerial and latent roots (morphological roots) in stems of some of the woody plants listed in table 1

Plant	Reference	Site
<i>Cotoneaster dammeri</i>	35	Parenchyma cells of divided leaf and bud gaps.
<i>Ficus carica</i>	13	Leaf traces, leaf gaps and medullary rays.
<i>Hedera helix</i>	36	Between the bark and vascular bundles.
<i>Lonicera japonica</i>	27	From cells between the cambium and pericycle at both nodes and internodes.
<i>Populus</i> Section <i>Aegeiros</i>	4	External to the cambium but in front of primary rays.
<i>Populus nigra italica</i> , <i>P. n. thevestina</i> , <i>P. simonii</i> , <i>P. trichocarpa</i>	37	Opposite medullary rays.
<i>Ribes alpinum</i>	37	Opposite medullary rays.
<i>Salix alba chermenisa</i> , <i>S. a. sericea</i> , <i>S. amygdalina</i> ,	37	Parenchyma of leaf gaps at the nodes; medullary rays of internodes.
<i>S. discolor</i>		
<i>Salix fragilis</i>	6,7	Outside the cambium in rays associated with leaf and branch gaps.
<i>Salix myrsinifolia</i>	37	Parenchyma of leaf gaps at the nodes; medullary rays of internodes.

ships between density or continuity of sclerenchyma and rooting potential in woody stem cuttings. They consider differences in ease of rooting to be related to the ease with which root initials are formed, not to the restriction of developing root primordia by sclerenchyma.

To end what I have to say this morning, I will briefly describe a few illustrations to give you an idea of how wound-induced roots arise in stems of *Hedera helix*. Figure 7 A, B shows cells at the outermost end of a ray, phloem ray parenchyma, with dense cytoplasm and deeply stained nuclei. The two large cells are the product of a recent anticlinal division. Groups of fibers (F) are present external and opposite the vascular bundles (VB). In figure 8 A, B a young root initial (RI) made up of a slightly organized group of cells is seen developing between two phloem traces. Fibers (F) flank the outer portion of the vascular bundles but they do not hinder the development of the young root. In a few cases (Fig. 9 A, B), vascular rays abut at their distal end on perivascular fibers. Examples of this kind are found even in easy-to-root stems of the juvenile growth phase. The rays often stain densely as shown in this

illustration. As the root initials continue to develop, cells divide and differentiate and begin to form organized tissues typical of root primordia (see fig. 10 A, B and 11). Vascular strands quickly differentiate into the actively growing primordia. Phloem elements of at least two vascular bundles and three rays take part in the formation of each root primordium. Scattered groups of fibers are unable to stop the last few stages of adventitious root formation (fig. 12). Roots with rootcaps

Table 3. Origin of wound-induced roots in stems of woody plants

Plant	Reference	Site
<i>Abelia grandiflora</i>	37	Callus.
<i>Abies</i> spp.	28	Callus.
<i>Acanthopanax spinosa</i>	37	Callus.
<i>Acanthus montanus</i>	34	Near vascular cambium.
<i>Caragana arborescens</i>	33	Secondary phloem; callus.
<i>Chamaecyparis</i> spp.	28	Vascular rays.
<i>Clematis</i> spp.	30	Vascular cambium.
<i>Cryptomeria</i> spp.	28	Vascular rays.
<i>Cupressus</i> spp.	28	Vascular rays.
<i>Forsythia suspensa</i>	33	Leaf and bud traces.
<i>Hedera helix</i> (juvenile)	15	Phloem ray parenchyma.
<i>Hedera helix</i> (mature)	16	Phloem ray parenchyma and callus.
<i>Ilex opaca</i>	21	Vascular cambium extended into callus; young phloem after auxin treatment.
<i>Larix</i> spp.	28	Bud traces.
<i>Ligustrum vulgare</i>	20	Outgrowths of lenticels.
<i>Picea</i> spp.	28	Leaf traces, callus.
<i>Pinus</i> spp.	28	Leaf traces, callus.
<i>Ribes alpinum</i>	33	Secondary xylem, cambium
<i>Rosa</i> 'Dorothy Perkins'	5	Secondary phloem.
<i>Rosa dilecta</i> 'Better Times'	31	Phloem ray parenchyma.
<i>Rubus idaeus</i>	32	Primary rays close to leaf traces.
<i>Rubus occidentalis</i>	32	From leaf and bud traces.
<i>Sambucus nigra</i>	20	Outgrowths of lenticels.
<i>Sciadopitys</i> spp.	28	Branch trees.
<i>Tamarix aphylla</i>	14	Outgrowths of lenticels.
<i>Taxus cuspidata</i>	19	Phloem ray cells and adjacent parenchyma away from basal cuts; secondary phloem near basal cuts.
<i>Thuja</i> spp.	28	Vascular rays.
<i>Thujopsis</i> spp.	28	Rays, leaf traces.
<i>Thujopsis dolobrata</i>	37	Callus.
<i>Ulmus campestris</i>	20	Outgrowths of lenticels.
<i>Vaccinium corymbosum</i>	23	Cambium and phloem.



present emerge above the basal cuts almost at right angle to the main axis of the stems. In cuttings made from stems of the mature growth phase roots also form from cells of the callus tissue (fig. 13 A, B).






Figure 7 A, B. Two cells of phloem ray parenchyma are the product of a recent anticlinal division. These cells have dense cytoplasm and deeply stained nuclei. PI, pith; P, phloem; X, xylem; VB, vascular bundles; F, fibers; IC, interfascicular cambium; FC, fascicular cambium.



Figures 8 A, B. A young root initial made up of a slightly organized group of cells is seen developing between two phloem traces. X, xylem; P, phloem; RI, root initial; F, fibers; C, cortex.





Figure 9 A, B. A densely stained vascular ray abuts at its distal end on a group of fibers. X, xylem; P, phloem; F, fibers; C, cortex; FC, fascicular cambium; IC, interfascicular cambium.




Figure 10 A, B. A young root primordium with well organized tissues. Phloem elements of at least two vascular bundles and three rays take part in the formation of a root primordium. PI, pith; XR, xylem ray; F, fibers; C, cortex. From: Girouard, R. M. 1967. *Canad. J. Bot.* 45:1877-1881.






Figure 11. Longitudinal section of an adventitious root primordium at a stage similar to or slightly younger than that shown in figure 11 A.

Figure 12. Transverse section of a stem with an adventitious root developing through cortical and epidermal tissues. Groups of fibers are unable to stop the last few stages of root formation. From: Girouard, R. M. 1967. *Canad. J. Bot.* 45:1883-1886.



(A)



(B)

Figure 13 A, B. In stems taken from mature phase plants adventitious roots also form in callus tissue at the basal end of cuttings. X<sup>1</sup>, primary xylem; X<sup>2</sup>, secondary xylem; P, phloem; C, cortex; E, epidermis; V, vessels; CA, callus; AM, apical meristem.

## LITERATURE CITED

- 1 Beakbane, A B 1961 Structure of the plant in relation to adventitious rooting *Nature* 192:954-955
- 2 Bloomberg, W J 1959 Root formation of black cottonwood cuttings in relation to region of parent shoot *For Chron* 35:13-18
- 3 Bloomberg, W J 1963 The significance of initial adventitious roots in poplar cuttings and the effect of certain factors on their development. *For Chron* 39 279-289
- 4 Braun, H J. 1963 Wurzelkeime in jungen und alteren Spriebachsen von *Populus* *Z Bot* 51:441-451
- 5 Carlson, M C 1933 Comparative anatomical studies of Dorothy Perkins and American Pillar roses I Anatomy of canes II. Origin and development of adventitious roots in cuttings *Boyce Thompson Inst Contrib* 5 313-330
- 6 Carlson, M C 1938 The formation of nodal adventitious roots in *Salix cordata* *Am J Bot* 25 721-725
- 7 Carlson, M C 1950 Nodal adventitious roots in the willow stems of different ages *Am J Bot* 37 555-561
- 8 Ciampi, C and Gellini, R 1958 Anatomical study on the relationship between structure and rooting capacity in olive cuttings *Nuovo giorn botan. ital* 65 417-424 (*Hort. Abstr* 30, *Abstr* 1197 1960)
- 9 Ciampi, C and Gellini, R 1963 The origin and development of adventitious roots in *Olea europaea*. the importance of anatomical structure in rootlet development. *Nuovo giorn botan. ital* 70 62-74 (*Hort Abstr* 35, *Abstr* 4422 1965).
- 10 Duhamel du Monceau, H L 1758 *La physique des arbres* 2 volumes H L Guerin and L F Delatou, Paris Plate XV, vol 2
- 11 Esau, K. 1960 *Anatomy of seed plants* John Wiley & Sons, Inc, New York 376 pp
- 12 Esau, K 1965. *Plant anatomy* 2nd ed John Wiley & Sons, Inc, New York. 767 pp
- 13 Gellini, R 1964 Studio sulla radicazione di talee di *Ficus carica* L. Osservazioni sulla struttura e sullo sviluppo di radici preformate *In Atti delle giornate di studio su la propagazione delle specie legnose* Università di Pisa, Istituto di coltivazioni arboree, pp 198-219
- 14 Ginzburg, C 1967 Organization of the adventitious root apex in *Tamarix aphylla* *Am J Bot* 54:4-8
- 15 Grouard, R M. 1967 Initiation and development of adventitious roots in stem cuttings of *Hedera helix* Anatomical studies of the juvenile growth phase *Canad J. Bot* 45:1877-1881.
- 16 Grouard, R M 1967 Initiation and development of adventitious roots in stem cuttings of *Hedera helix*. Anatomical studies of the mature growth phase. *Canad J Bot* 45 1883-1886
- 17 Goodin, J R 1965 Anatomical changes associated with juvenile-to-mature growth phase transition in *Hedera* *Nature* 208 504-505.
- 18 Hartmann, H T. and Kester, D E. 1959. *Plant propagation — principles and practices* Prentice-Hall Inc, Englewood Cliffs, N. J 559 pp
- 19 Hiller, C. H 151 A study of the origin and development of callus and root primordia of *Taxus cuspidata* with reference to the effects of growth regulator Master's thesis Cornell Univ, Ithaca, N Y 68 pp
- 20 Jeanmaire, L and Gorenflot, R 1964 L'hypertrophie des lenticelles sous l'influence de l'eau *C R Acad Sci* 258 3537-3540
- 21 Klinck, D S 1960 The origin of adventitious roots on the stem cuttings of *Ilex opaca* as influenced by wounding and synthetic growth substances. Master's thesis Ohio State University 56 pp
22. Mahlstedt, J P. and Haber, E S 1957. *Plant propagation* John Wiley & Sons, Inc, New York 413 pp
- 23 Mahlstedt, J. P and Watson 1952 An anatomical study of adventitious root development in stems of *Vaccinium corymbosum* *Bot Gaz* 113 279-285.
- 24 Reed, H S 1942 A short history of the plant sciences. *Chronica Botanica* Co, Waltham, Mass 320 pp
- 25 Ruggeri, C 1964 Osservazioni anatomo-fisiologiche sul radicamento di alcuni cloni di pioppo *In Atti delle giornate di studio su la propagazione*

- delle specie legnose. Università di Pisa, Istituto di coltivazioni arboree pp 254-266
- 26 Sachs, R. M., Loreti, F., and De Bie, J. 1964. Plant rooting studies indicate sclerenchyma tissue is not a restricting factor. *Calif Agric* 18 (No. 9) 4-5
  - 27 Sandison, S. 1934. Rooting of cuttings of *Lonicera japonica*. A preliminary account. *New Phytol* 33 211-217
  - 28 Satoo, S. 1956. Anatomical studies on the rooting of cuttings in coniferous species. *Bull Tokyo Univ Forests* No. 51, p. 109-158 (For Abstr. 19, Abstr. 1164, 1958)
  - 29 Shapiro, S. 1958. The role of light on root primordia in the stems of the Lombardy poplar. In *The physiology of forest trees* (Ed. K. Thimann). The Ronald Press, New York p. 445-465
  - 30 Smith, E. P. 1928. A comparative study of the stem structure of the genus *Clematis* with special reference to anatomical changes induced by vegetative propagation. *Roy Soc Edinburgh, Trans* 55 643-664
  - 31 Stangler, B. 1956. Origin and development of adventitious roots in stem cuttings of chrysanthemum, carnation and rose. *Cornell Agr. Exp. Sta. Mem.* 342, 24 pp
  - 32 Sudds, R. H. 1935. The origin of roots in several types of red and black raspberry stem cuttings. *Proc Amer. Soc. Hort. Sci.* 33 380-385
  - 33 Swartley, J. C. 1942. Adventitious root initiation in *Forsythia*, *Ribes* and *Caragana*. Master's thesis, Ohio State Univ., Columbus, Ohio
  - 34 Taylor, G. 1926. The origin of adventitious growths in *Acanthus montanus*. *Bot. Soc. Edinburgh, Trans. and Proc.* 29 291-296
  - 35 Tomlinson, P. B. 1961. *Anatomy of the monocotyledons. II. Palmae*. Oxford Univ. Press. 453 pp
  - 36 Trecul, G. 1846. Recherches sur l'origine des racines. *Ann. Sci. Nat. Bot.* ser. 3, 6 340-350
  - 37 van der Lek, H. A. A. 1924. Over de wortelvorming van houtige stekken (English summary). *Meded. Landbouwhoogeschool, Wageningen* 28 1-230
  - 38 Wolfe, F. H. 1931. The origin of adventitious roots in *Cotoneaster dammeri*. Master's thesis. Northwestern Univ., Evanston, Illinois. 15 pp

MODERATOR CANNON: Our next speaker is also from Canada. Mr. Joerg Leiss will speak on outdoor grafting.